



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

with coarser appressed hairs in all our specimens. It must be that the first description is wrong on this point. Then, the leaves are in no case "7—8 lines wide," but $\frac{3}{8}$ in. to $\frac{1}{2}$ in.; the Angeles Bay plant having some leaves as wide as $\frac{3}{8}$ in. Furthermore, the description has "Calyx 4-parted . . . Stamens 6 according to Bonpland. . . . Achene lenticular." I have frequently found the calyx 5-parted, stamens as many as 8, and the achenes in the Angeles Bay and the Orizaba plant—the latter from one of the stations cited in Biol. Centr. Am. l. c.—as well as in the Texas plant, are triangular: all, however, of the same size, and all "umbonate," as in the first description.

This species is distinguished from *P. persicaria* L. by its narrower, longer leaves, more slender spikes and smaller achenes.—J. M. HOLZINGER, *Department of Agriculture, Washington, D. C.*

New Mosses of North America.—The following brief diagnoses are published in advance of fuller descriptions in order to secure priority. We hope to prepare shortly the fifth number of our series under the above title, in THE BOTANICAL GAZETTE.

Dichodontium olympicum n. sp.—A *D. pellucida* jam multo robustiore primo visu differt: foliis valde papillosis, toto fere ambitu minute denticulatis, capsulaque basi strumosa. Planta humilis, vix 1 cent. alta.—Olympic Mts., Wash. (*L. F. Henderson*.)

Grimmia Hendersoni n. sp.—*G. decipiente* Lindb. (*G. Schultzii* Wils.) proxima sed ab ea pedicello longiore, capsula subcylindrica magis elongata, operculo longius rostrato et foliorum rete basilari multo laxiore facillima distinguenda.—Hood River, Oregon, on dry rocks. (*L. F. Henderson*).

Eucalypta lacera n. sp.—Ab *Enc. vulgari* proxima differt calyptra basi lacerata, peristomio e membrana alba fugacissima lacerata composito et pedicello longiore.—Milwaukie, Oregon, Willamette River, (*L. F. Henderson*).

Leskea obtusa n. sp.—Formis robustioribus *L. polycarpæ* similis. sed ab illis primo aspectu foliis obtusis marginibus planis distincta, A *L. obscura* habitu valde robustiore, foliis majoribus et magis obtusis, costa brevior et capsula majore et longiore quoque differt.—Bethlehem, Pa., mixed with *Anomodon obtusifolius* (Rau.) Chinchuba, La., near Mandeville, on trees (*Langlois*).—F. RENAULD AND J. CARDOT, *Monaco, and Stenay, France*.